

Ribosomal DNA sequence analysis of *Heterodera filipjevi* and *H. latipons* isolates from Russia and comparisons with other nematode isolates

Virginia R. Ferris*, Sergei A. Subbotin**, Anita Ireholm***, Yitzhak Spiegel****, Jamal Faghihi* and John M. Ferris*

*Department of Entomology, Purdue University, West Lafayette, IN, 47907-1158, U.S.A., e-mail: vferris@entm.purdue.edu

**Institute of Parasitology of Russian Academy of Sciences, Leninskii Prospect, 33, Moscow, 117071, Russia,

***Swedish University of Agricultural Science, S-230 53 Alnarp, Sweden,

****Division of Nematology, Agricultural Research Organization, The Volcani Center, P. O. Box 6 Bet Dagan, Israel 50250.

Accepted for publication 6 August 1999

Summary. Ribosomal DNA (ITS rDNA) sequence data for two geographically separated isolates of *Heterodera filipjevi* from Russia were identical, and were 99.7% similar to rDNA sequence from Swedish East and West Gotland strains of *H. avenae*. These data are consistent with the view that the 'Gotland strain' of *H. avenae* and *H. filipjevi* are conspecific. The ITS rDNA sequence from an isolate of *H. latipons* from the Rostov region of Russia was 98.1% similar to the ITS rDNA sequence of an isolate from Gilat, Israel, the type locality of *H. latipons*, which suggests the possibility that the two morphologically similar, but geographically separated, isolates might be sibling species.

Key words: *Heterodera avenae*, *H. filipjevi*, *H. latipons*, ribosomal DNA, rDNA sequence, phylogenetic analysis, Russia, Israel, Sweden.

Heterodera filipjevi (Madzhidov, 1981) Stelter, 1984 is an important nematode pest of cereals in European and Middle Asian countries (Subbotin, *et al.*, 1999). It was believed that this species, described from Tadzhikistan, had a limited geographic distribution, but recent morphological, biochemical and molecular studies indicate that populations from widely dispersed geographic regions, originally identified as *H. avenae* Wollenweber, 1924, or strains of *H. avenae* (Gotland and pathotype 3), or even as *H. mani* (Mathews, 1971), are probably conspecific with *H. filipjevi*. Ferris *et al.* (1989, 1994) compared isolates of *H. avenae sensu stricto* from Sweden, Australia and America with isolates of the Swedish 'Gotland strain' of *H. avenae* using 2-D PAGE protein patterns and sequence data for the two internal transcribed spacers of the ribosomal RNA gene (ITS rDNA). These researchers found significant differences between the 'Gotland strain' isolates of *H. avenae* and *H. avenae sensu stricto*. Subbotin *et al.* (1999) reported that, based on canonical discriminant analysis of morphometric and restriction fragment length polymorphism (RFLP) data, the East Gotland strain from Etelhem Sweden was similar/identical to *H. filipjevi* populations from Russia.

An objective of the present research was to compare sequence data for ITS rDNA from two populations of *H. filipjevi* from Russia with previously obtained ITS rDNA sequence data from 'Gotland strain' isolates of *H. avenae*, including an Etelhem isolate (Ferris *et al.*, 1994).

Heterodera latipons Franklin, 1969 is an important cereal cyst nematode in many Mediterranean countries (Franklin, 1969). Widespread geographic populations have many morphological similarities, but the extent of genetic diversity among these populations is presently unknown. It is possible that some of these populations might actually comprise sibling species. A second objective of this study was to compare ITS rDNA sequence from a Russian isolate of *H. latipons* with ITS rDNA sequence for *H. latipons* from the type locality in Israel.

MATERIALS AND METHODS

Nematode isolates. Populations used for this study included two isolates of *H. filipjevi* and one of *H. latipons* from Russia. The *H. filipjevi* isolates were from Pushkin, Leningrad region (host: false wheat - *Elytrigia repens* L.) and from Vad, Nizhnii Novgorod

region (host: oats - *Avena sativa* L.). The *H. latipons* isolate (host: false wheat - *Elytrigia repens* L.) was from the Rostov region. Other isolates from earlier work (Ferris *et al.*, 1989, 1994) used for comparisons in this study, included the following isolates from Sweden: *H. avenae sensu stricto* of pathotype Ha12 from Nässja, three isolates of the East Gotland pathotype (from Etelhem, Stacketorp and Alnarp), and two isolates of the 'West Gotland' pathotype (from Hjelmsäter and N. Härene). All cereal cyst nematodes used in the earlier studies were from growth chamber cultures of barley cv. Varde (*Hordeum vulgare* L.). Details can be found in the original publications (Ferris *et al.*, 1989, 1994). The isolate of *H. latipons* from oats growing in Israel was from Gilat, Israel, the type locality (Franklin, 1969).

Amplification, cloning and sequencing. The methods for handling the nematodes and obtaining the rDNA were essentially as previously described by Ferris *et al.* (1993, 1994, 1995), and are briefly summarized here. Female nematodes (usually one or two) taken from a vial of 70% alcohol were rinsed in sterile water and crushed in 20 µl cold TE buffer, using a Radnoti (Thomas Scientific, Swedesboro, NJ, USA) 25 µl-size glass homogenizer. The homogenate was either used immediately or stored at -20° C. Five such preparations were usually made for each isolate. Prior to amplification by polymerase chain reaction (PCR), the homogenate was thawed briefly, and spun in a microfuge for 3 minutes at 16,000 g. About 15 µl of the supernatant was discarded, 60 µl Instagene (BioRad, Hercules, CA, USA) was added to the pellet, and the procedure completed according to manufacturer's directions for use of Instagene. Usually, 10 µl of a 1:10 dilution of the preparation was used for each 25 µl PCR reaction, but occasionally this was varied to improve amplification. Standard PCR (Saiki, 1990), as described in Ferris *et al.* (1993) was used with reagents from Perkin Elmer (Norwalk, CT, USA) and Promega (Madison, WI, USA), and a COY Tempcycler model 50 (COY, Inc., Ann Arbor, MI, USA). The PCR was for 30 cycles with denaturation at 94 °C for 1 min, annealing at 43 °C for 1 min, and extension at 72 °C for 1 min. The amplified DNA band was monitored by electrophoresis in SeaPlaque GTG agarose (FMC, Rockland, ME, USA), excised, and diluted in 1 ml sterile water. Thereafter, approximately 1 ng of the diluted DNA was amplified for 30 cycles as described previously, except that annealing was at 60 °C. Primers for PCR amplification were those described by Ferris *et al.* (1993). The amplified region of about 1 kb in length spanned the two ITS regions and included the 5.8S gene that lies between them. Multiple amplifications were carried out for each

isolate in order to be able to detect any sequence variation, but none occurred with these isolates. Amplified rDNA was cloned into the pGEM-T vector (Promega) and transformed into the *E. coli* strain JM109 according to the manufacturer's protocols. Clones were checked, using PCR, for the correct insert of the predicted size (approximately 1kb), and plasmid preparations made using the Wizard Plus Miniprep kit (Promega). Manual sequencing, using Sequenase version 2.0 (U.S. Biochemical, Cleveland, OH, USA), was used for all earlier isolates previously described (Ferris *et al.*, 1994); but sequences were obtained for the isolates from Russia with an automatic sequencing system (ALFexpress, Pharmacia Biotech, Piscataway, NJ, USA) used in the Purdue Sequencing Laboratory. In each instance, the double-stranded sequencing was from multiple clones, usually 2 to 4 clones were sequenced per isolate, to assure accuracy. Internal primers used for manual sequencing were those reported previously by Ferris *et al.* (1994). Sequence obtained from the automatic system was long enough, and of sufficient quality, that internal primers were not necessary.

Sequence comparisons. Sequence data for each isolate, including ITS1, 5.8S, and ITS2 rDNA, were compared with the data obtained for the other isolates, using the computer program GAP in the Sequence Analysis Software Package of the Genetics Computer Group, followed by PILEUP (Devereaux *et al.*, 1984).

RESULTS

Sequence data for the Pushkin isolate of *H. filipjevi*, the Etelhem isolate of the East Gotland strain, and the Russia and Israel isolates of *H. latipons*, are shown in Fig. 1. The rDNA sequences of the two isolates of *H. filipjevi* were identical. The Swedish East and West Gotland sequences also were identical, and differed from the sequence of the *H. filipjevi* isolates by only three base pairs (bp 952-954 in ITS2, Fig. 1). Pairwise percentage sequence similarities for these isolates, and for *H. avenae*, are shown in Table 1. The *H. filipjevi* sequence was 99.7% identical to the sequence from the Swedish East and West Gotland strains. The rDNA sequence of the isolate of *H. latipons* from Russia was 98.1% similar to that from the type locality, Gilat, Israel.

DISCUSSION

Molecular and biochemical data have contributed to improvements in the identification and placement of nematode populations of species of the *H. avenae* group. In addition to the research described above, Bossis and Rivoal (1996), using 2-D PAGE,

	1				50					100
FIL	CCCAAGTGAT	TCCATTACACC	ATCTACCTGT	GCTGTCCAGT	TGGAACGTTG	CTTGGCACCA	CCACATGCCC	CCGCTCTGCTG	TTGAGACGGG	CACGGACCGG
GOT
LAT-RT.GT	G.C.....T.T.T.A
LAT-IT.GT	G.C.....T.T.T.T.....
	101				150					200
FIL	TCGAGATGGT	CTGTGGGCAC	GGGACAACAC	TGAGCGGACG	GC-TACCCTG	CGAGCACTCT	GTGCTTTGGG	GTGTTCTCCG	TCGATGGTGC	TTGGTATACT
GOT
LAT-RA.T.CA.....AT.A.T.AA	---G.T.GAG	C.....C.....
LAT-IA.A.....AT.A.T.AA	---G.T.GAG	C.....C.....
	201				250					300
FIL	GACTCGTTGC	TGAGCAAAGT	GATAATAAAA	GCCTGAGGTT	TG--GCTGCG	AGGCAATTGA	GTTGGTGGCG	GACCGCTCGC	TGGTGGGTTT	GCTGTACCGT
GOT
LAT-RG.G.G---T.A.T.C.....AA
LAT-IG.G.G.CT.T.A.T.C.....AA
	301				350					400
FIL	CGAAGGGAGC	ACGCTCGTTG	GGTAACCCAA	CGGCTGTGCT	GGTGTCTGTG	CGTCGTTGAG	CGGTTGTTGT	GCCTGGCACA	TGTGACACAC	TGGCCTGGGA
GOT
LAT-RG.A.C.AA---A.A.....A.
LAT-IG.A.C.AA---A.A.....A.
	401				450					500
FIL	AGGTGGTTCT	-TTCCTGGCC	TTACGAACCG	TAAGTACCGG	TGTGTCTGTG	CTTGTGTGCT	ACGTCCGTGG	CCGTGATGAG	ACGACGTGGT	AGGGTCCGTG
GOT
LAT-R	TAT.....G.	G.....T.	G.....TG.A.C.C.....
LAT-I	TAT.....G.	G.....T.	TG.....	TA.....G.C.C.....
	501				550					600
FIL	CTCTGCTTAT	AGCACGTGGC	TTAAGACTTA	ATGAGTGTCA	GCTAGGCACC	GCCAGTGTTT	TTTTTCATTT	A--TTTTTTT	GACCACTTCT	TTGTTGAAGG
GOT
LAT-RT.G.C.C.--T.A.A.....C.....A
LAT-IT.G.C.C.TTT.A.A.....C.....A
	601				650					700
FIL	AAGAAATTC	AGTCTTATCG	GTGGATCACT	CGGCTCGTGG	ATCGATGAAG	AACGCAGCCA	ACTGGCATAA	TTAGTGCAGAA	CTGCAGAAAC	CTTGAACACA
GOT
LAT-RT.
LAT-I
	701				750					800
FIL	AAACTTTTGA	ATGCACATGG	CGCCATTGGA	GTTACATCCA	TTGGCACGCC	TGGTTCAGGG	TCGTTATCAT	AAAA-GCACT	GCTGTGCCTG	TTATGTTGGT
GOT
LAT-RT.G.
LAT-IC.G.
	801				850					900
FIL	GAGATCATGT	CTGCAGGACG	TGTTCTTGGC	CTATTCTTGA	AAATGCTCGG	CCGTGGAGTG	TGGTTGTGTT	GGCGCGAAAC	TGTCAGGTTT	TITCGCGTTT
GOT
LAT-RTT.TCATT	G.....C.T.T.C.A.....TTTGCG..
LAT-ITT.TCATT	G.....C.T.T.C.A.....TTCGCG..
	901				950					982
FIL	TACGGTCCGT	AACTAGGACG	CAACTGCTCT	CCATGTGTGC	T--GGGTGGA	ATGTTGCTG	GTAGGCATTC	CGTGTTTGAA	TT	
GOTCTG.	
LAT-RA.T.G.G.T.A.TG..CG.A.C.TCA.	
LAT-IA.T.G.T.A.TG..CG.A.CG.TCT.	

Fig. 1. Nucleotide bases of rDNA sequence of ITS1, 5.8S (italics), and ITS2 for *Heterodera filipjevi* from Russia (FIL); *H. avenae*, 'East Gotland' strain from Etelhem, Sweden (GOT); *H. latipons* from Russia (LAT-R); and *H. latipons* from Israel (LAT-I). All base notations are for the non-transcribed strand. Periods indicate sequence identity and hyphens indicate gaps. Sequence differences are uncorrected for multiple changes at a site.

also showed clear genetic dissimilarities between *H. avenae sensu stricto* and the 'Gotland strain' isolates. López-Braña *et al.* (1996) used random amplified polymorphic DNA (RAPD) patterns to separate *H. avenae sensu stricto* from Gotland strain populations from Spain (Torralba de Calatrava) and Sweden (Etelhem).

Clarification of the taxonomy of strains and populations of cereal cyst nematodes has been progressed through comparison of populations from the type locality of *H. filipjevi* with other nominal *H. avenae* and 'Gotland strain' populations. Subbotin *et al.* (1996, 1999) showed that cereal cyst nematode populations from Russia and the Ukraine, previously assigned to *H. avenae*, were morphologically and

morphometrically conspecific with the Tadzhikistan population of *H. filipjevi*. The authors confirmed these findings with biochemical data from isoelectric focusing patterns of protein extracts. They further showed that only *H. filipjevi* is widely distributed in the European part of Russia. On the basis of electrophoretic and morphological studies, Rumpfenhorst *et al.* (1996), Sturhan (1996) and Sturhan and Rumpfenhorst (1996) reported that most of the cereal cyst nematode populations from Turkey, Iran and Germany, previously identified as *H. avenae* or *H. mani*, were identical to *H. filipjevi*. Sturhan and Rumpfenhorst (1996) published identical protein patterns from cysts of the 'Gotland strain', the strain known as pathotype 3, and *H. filipjevi* populations.

Table 1. Pairwise percentage similarity values for the nucleotide bases shown on Figure 1 plus comparable rDNA data for *H. avenae* (Ferris *et al.*, 1994)*.

	FIL	GOT	LAT-R	LAT-I	HAV
FIL	-	99.7	90.2	89.4	96.5
GOT	-	-	90.0	89.5	96.8
LAT-R	-	-	-	98.1	87.0
LAT-I	-	-	-	-	87.1

*Abbreviations for species/isolates are as in Figure 1. HAV = *H. avenae*.

Bekal *et al.* (1997), in a study in which internal transcribed spacer (ITS) rDNA was cut with restriction enzymes, found no differences in restriction fragment length polymorphisms (RFLPs) between *H. filipjevi* populations and 'Gotland strain' populations from Sweden (Etelhem), Spain (Torralba de Calatrava) and Bulgaria. Subbotin *et al.* (1999) found, using morphometric and rDNA - RFLP data, that the populations used by Bekal *et al.* (1997) from Etelhem, Sweden and from Spain were conspecific with further populations of *H. filipjevi* from Russia.

Data from our research are consistent with suggestions of Sturhan & Rumpfenhorst (1996), Bekal *et al.* (1997), and Subbotin *et al.* (1999) that the Swedish Gotland strain isolates are conspecific with *H. filipjevi*; and on the basis of presently available data we believe them to be so. Similarity in ITS rDNA between the *H. latipons* isolate from Russia and that from Israel is high (98.1%), suggesting a close phylogenetic relationship. However, similarities in ITS rDNA sequence between *Globodera rostochiensis* and *G. pallida*, the two potato cyst nematode species, are only slightly less (96.7%) than for the two *H. latipons* populations (Ferris *et al.*, 1995). Our original diagnosis that the Russian isolate used here was *H. latipons* was based on data discussed in Subbotin *et al.* (1999). More recent observations by one of us (SAS) of further populations from Asia, and comparisons with type slides from the Rothamsted collection, suggest the possibility that our isolate from Russia, could, in fact, be a sibling species of *H. latipons*. In addition to further morphometric and other research on similar populations from diverse parts of the world, molecular sequence data based on other genes will likely be needed to resolve the question of the extent of variability and species integrity of these geographic isolates that appear to be similar to *H. latipons*.

ACKNOWLEDGEMENTS

Supported in part by National Science Foundation grant DEB9705740 to VRF.

REFERENCES

- Bekal, S., Gauthier, J.P. & Rivoal, R. 1997. Genetic diversity among a complex of cereal cyst nematodes inferred from RFLP analysis of the ribosomal internal transcribed spacer region. *Genome* 40: 479-486.
- Bossis, M. & Rivoal, R. 1996. Protein variability in cereal cyst nematodes from different geographic regions assessed by two-dimensional gel electrophoresis. *Fundamental and Applied Nematology* 19: 25-34.
- Devereaux, J.R., Haeberli, P. & Smithies, O. 1984. A comprehensive set of sequence analysis programs for the VAX. *Nucleic Acids Research* 12: 387-395.
- Ferris, V.R., Faghihi, J., Ireholm, A. & Ferris, J.M. 1989. Two dimensional protein patterns of cereal cyst nematodes. *Phytopathology* 79: 927-933.
- Ferris, V.R., Ferris, J.M. & Faghihi, J. 1993. Variation in spacer ribosomal DNA in some cyst-forming species of plant parasitic nematodes. *Fundamental and Applied Nematology* 16: 177-184.
- Ferris, V.R., Ferris, J.M., Faghihi, J. & Ireholm, A. 1994. Comparisons of isolates of *Heterodera avenae* using 2-D PAGE protein patterns and ribosomal DNA. *Journal of Nematology* 26: 144-151.
- Ferris, V.R., Miller, L.I., Faghihi, J. & Ferris, J.M. 1995. Ribosomal DNA comparisons of *Globodera* from two continents. *Journal of Nematology* 27: 273-283.
- Franklin, M.T. 1969. *Heterodera latipons* n. sp., a cereal cyst nematode from the Mediterranean region. *Nematologica* 15: 535-542.
- López-Braña, I., Romero, M.D. & Delibes, A. 1996. Analysis of *Heterodera avenae* populations by the random amplified polymorphic DNA technique. *Genome* 39: 118-122.
- Rumpfenhorst, H. J., Elekçioğlu, I.H., Sturhan, D., Öztürk, G. & Enneli, S. 1996. The cereal cyst nematode, *Heterodera filipjevi* (Madzhidov) in Turkey. *Nematologia mediterranea* 24: 135-138.
- Saiki, R.K. 1990. Amplification of genomic DNA. In: *PCR Protocols* (M. A. Innis, D. H. Gelfand, J. J. Sinski, & T. J. White, Eds.) pp. 13-20. San Diego, CA: Academic Press.

- Sturhan, D. 1996.** Occurrence of *Heterodera filipjevi* (Madzhidov, 1981) Stelter, 1984 in Iran. *Pakistan Journal of Nematology* 14: 89-93.
- Sturhan, D. & Rumpfenhorst, H.J. 1996.** Untersuchungen über den *Heterodera avenae*-Komplex. *Mitteilungen aus der Biologischen Bundesanstalt für Land-Forswirtschaft, Berlin-Dahlem*, 317: 75-91.
- Subbotin, S.A., Rumpfenhorst, H.J. & Sturhan, D. 1996.** Morphological and electrophoretic studies on populations of the *Heterodera avenae* complex from the former USSR. *Russian Journal of Nematology* 4: 29-38.
- Subbotin, S.A., Waeyenberge, L., Molokanova, I.A. & Moens, M. 1999.** Identification of *Heterodera avenae* group species by morphometrics and rDNA-RFLPs. *Nematology* 1: 195-207.
-

Ferris V.R., Subbotin S.A., Ireholm A., Spiegel Y., Faghihi J., Ferris J.M. Анализ последовательностей рибосомальной ДНК изолятов *Heterodera filipjevi* и *H. latipons* из России и сравнение их с другими изолятами нематод.

Резюме. Последовательности рибосомальной ДНК (ITS rDNA) двух географически удаленных изолятов *Heterodera filipjevi* из России оказались идентичными и на 99.7% соответствовали рДНК последовательностям Восточной и Западной Готландской расы *H. avenae* из Швеции. Полученные данные подтверждают, что Готландская раса *H. avenae* и *H. filipjevi* конспецифичны. Последовательности рибосомальной ДНК изолята *H. latipons* из Ростовской области России были на 98.1% сходны с последовательностями ITS рДНК изолята этого вида из Гилата (Израиль) - типового места обнаружения этого вида. Эти данные показывают, что два этих морфологически сходных и географически удаленных друг от друга изолята возможно представляют собой виды-близнецы.
